

Poster presentation

Therapeutic potential of CNP for skeletal dysplasias

Akihiro Yasoda*¹, Hidetomo Kitamura², Toshihito Fujii¹, Eri Kondo¹, Naoaki Murao³, Masako Miura¹, Naotetsu Kanamoto¹, Yasato Komatsu¹, Hiroshi Arai¹ and Kazuwa Nakao¹

Address: ¹Department of Medicine and Clinical Science, Kyoto University Graduate School of Medicine, Kyoto, Japan, ²Project Management Department, Chugai Pharmaceutical Co., Ltd., Tokyo, Japan and ³Fuji-Gotemba Research Laboratories, Chugai Pharmaceutical Co., Ltd., Gotemba, Shizuoka, Japan

Email: Akihiro Yasoda* - yasoda@kuhp.kyoto-u.ac.jp

* Corresponding author

from 4th International Conference of cGMP Generators, Effectors and Therapeutic Implications Regensburg, Germany. 19–21 June 2009

Published: 11 August 2009

BMC Pharmacology 2009, 9(Suppl 1):P74 doi:10.1186/1471-2210-9-S1-P74

This abstract is available from: <http://www.biomedcentral.com/1471-2210/9/S1/P74>

© 2009 Yasoda et al; licensee BioMed Central Ltd.

Background

We had revealed that C-type natriuretic peptide (CNP) is a potent stimulator of endochondral bone growth by using transgenic and knockout mice: transgenic mice with targeted overexpression of CNP in cartilage develop prominent skeletal overgrowth [1], whereas mice depleted with CNP exhibit impaired endochondral bone growth [2]. We planned to translate this effect of CNP into the therapy for skeletal dysplasias, of which the effective drug therapy is not established yet. Here we investigated the effect of CNP on the impaired bone growth of mice model of achondroplasia, the most common form of skeletal dysplasias.

Results

First, we investigated the effects of plasma CNP on impaired bone growth in achondroplastic mice that specifically overexpress CNP in the liver under the control of human serum amyloid P component promoter. Increased plasma CNP stimulated the impaired growth of achondroplastic bones, and they grew almost comparable to those of wild type mice at the age of 6 weeks. Then we treated achondroplastic mice with continuous intravenous infusion of CNP at the dose of 1 µg/kg/min, and successfully rescued the shortness of most bones formed endochondral ossification at the end of 3-week experimental period. Figures 1 and 2.

Conclusion

Our results indicate that treatment with CNP is a potential therapeutic strategy for skeletal dysplasias, including achondroplasia, in humans.




Figure 1
Soft x-ray picture of wild type mice (Wt), achondroplastic model mice (Ach), and achondroplastic model mice treated with CNP at the dose of 1 µg/kg/min.




Figure 2
Lengths of bones measured on soft x-ray film.

Acknowledgements

We would like to thank Dr. David Ornitz (Washington University Medical School) for giving us mice model of achondroplasia.

References

1. Yasoda A, Komatsu Y, Chusho H, Miyazawa T, Ozasa A, Miura M, Kurihara T, Rogi T, Tanaka S, Suda M, Tamura N, Ogawa Y, Nakao K: **Overexpression of CNP in chondrocytes rescues achondroplasia through a MAPK-dependent pathway.** *Nat Med* 2004, **10**:80-86.
2. Chusho H, Tamura N, Ogawa Y, Yasoda A, Suda M, Miyazawa T, Nakamura K, Nakao K, Kurihara T, Komatsu Y, Itoh H, Tanaka K, Saito Y, Katsuki M, Nakao K: **Dwarfism and early death in mice lacking C-type natriuretic peptide.** *Proc Natl Acad Sci USA* 2001, **98**:4016-4021.

Publish with **BioMed Central** and every scientist can read your work free of charge

"BioMed Central will be the most significant development for disseminating the results of biomedical research in our lifetime."

Sir Paul Nurse, Cancer Research UK

Your research papers will be:

- available free of charge to the entire biomedical community
- peer reviewed and published immediately upon acceptance
- cited in PubMed and archived on PubMed Central
- yours — you keep the copyright

Submit your manuscript here:
http://www.biomedcentral.com/info/publishing_adv.asp

